

Family Foot & Ankle Care

Philip C. Caswell, D.P.M.

Diplomate, American Board of Podiatric Surgery

Board Certified in Foot Surgery Fellow, American College of Foot and Ankle Surgeons

> 122 North Church Road (Route 94), Sparta, NJ 07871 Phone (973) 300-9151 Fax (973) 300-9175

Date:		Patient:	
Ar	rea(s) of care:		
	POST-OPER	ATIVE INSTRUCTIONS FOR TOENAIL SURGERY	
1.	Remove the bandage tomorrow morning. If the bandage adheres to the surgical site, soak as directed with the bandage on. When the bandage loosens, gently remove from the surgical site.		
2.	Soak your toe(s) in one quart of warm water with 4 Tablespoons of Epsom salts. Soak your foot/feet for 15 – 20 minutes twice a day. Try to soak your foot/feet at the same time every day (once in the morning and once at night). It is acceptable to soak more frequently. *However, you must soak at least once a day.		
3.	Remove your toe(s) from the soak water. Cleanse the surgical site(s) with a clear cotton swab (Q-tip). Clean the surgical site(s) (the area of the nail that has been removed) with the cotton swab twice. Do not use the same side of the swab more than once. Do not use Hydrogen Peroxide on the surgical site.		
4.		cream (ex. Neosporin Cream) to the surgical site. Do not use interferes with normal healing.	
5.	Cover the surgical	site(s) with a Band-Aid.	
6.		ctions until otherwise notified by your surgeon. DO not perative care on your own.	
7.	foot, foul smelling surgeon at (201) 8	ver, chills, increasing discomfort, redness of the toe(s) and/or and/or thick drainage, and increased swelling please call your 74-3561. If you have Diabetes and experience an unexplained bood sugar, call your surgeon.	
8.	You have been give a. Pain Medica	en the following medication(s). Your instructions are:	
	b. Antibiotic:		
9.	It is normal to have some non-infected clear or watery yellow drainage for severa		

weeks after the surgery.